

Managing freshly calved dairy cows

THE transition period, defined as three weeks pre-calving (Figure 1) and three weeks post-calving, is the highest risk period for the dairy cow during the production cycle.

This article will discuss some of the factors that can improve performance through the post-calving transition period and also touches on metritis. The second article in the series will review the factors that can affect the pre-calving period.

The reduction in dry matter intake (DMI) during the immediate pre-partum period and the lag in its rise post-calving, together with the dramatic increase in energy demands as lactation commences, are well-documented¹. Over the same period, the requirement for calcium jumps from ~17g/d to ~50g/d².

This discrepancy between the increase in demands and DMI results in a period of negative energy balance for the modern Holstein cow during early lactation, and is a normal process. Modern cows are capable of mobilising body reserves (primarily in the form of

JIM WILLSHIRE

BVSc, DCHP, MRCVS

discusses minimising negative energy balance to avoid related problems in cows during the productive cycle by maximising a well-constructed feed ration

fats) to meet this shortfall and continue to yield, which, as DMI increases through lactation, are replenished.

The mobilisation of body reserves can be observed in the milking cow through:

- condition loss during early lactation measured through body condition scoring (BCS)
- increased circulating levels of non-esterified fatty acids (NEFA)
- increased circulating beta-hydroxybutyrate (BHB) levels

The increased energy and calcium demands mean presenting the correct ration and maximising DMI are essential to minimise the level of negative energy balance that occurs. Should the degree of negative energy balance or hypocalcaemia slip above a threshold then they can act as gateway diseases for:

- compromised immune systems³
- displaced abomasums⁴⁻⁷
- extended calving to conception period^{8,9}
- increased duration and severity of mastitis¹⁰
- increased prevalence of mastitis (clinical and subclinical)^{6,11,12}
- metritis¹³
- reduced yields¹⁴
- retained foetal membranes^{15,16}

The impact and risk of both subclinical ketosis and hypocalcaemia can be reduced by maximising the intake of a well-constructed ration.

After calving, the aim should be to move the cow from the calving area and ration to a higher energy and calcium density ration as quickly as possible. In larger herds, very often a fresh cow group is employed where the stock-




Figure 1. Cows pre-calving in cubicles.

ing density is kept as low as possible – fresh feed is always available and access to the parlour is convenient to try to ensure removal from feed is minimised. Feed space should be maximised and, ideally, approaching 1m/cow¹⁷ to try and minimise the effect negative interactions/guarding.

Social interactions

Due to the social nature of cows it has been suggested changes in groups can have negative effects on DMI as the group hierarchy is established. These social interactions can pose a significant risk to the freshly calved cow. However, the evidence as to whether these interactions have a negative effect on either DMI or energy balance is mixed^{18,19} and the potential for minimising the number of group changes given typical UK herd size is limited.

Sufficient time for comfortable rest should form a significant part of a dairy cow's daily time budget and insufficient or uncomfortable rest can not only have sig-

nificant impacts on DMI and, consequently, energy balance and milk production²⁰, but also on udder and foot health.

Depending on the herd, fresh cows may be provided with loose yards or cubicles. There are pros and cons to both, the details of which, however, are beyond the scope of this article.

Provided the system is well designed and managed there is little to decide between them. Suggested cubicle dimensions for mature Holstein cows are given in Figure 2 and areas for straw yards in Table 1.

The presence of concurrent disease will have negative effects on DMI, and its early detection and effective treatment is critical. In larger herds, a fresh cow programme is often instituted as a proxy for classic stockmanship. These programmes will often use a combination of routine urine ketone testing, rectal temperatures, milk yield and vaginal examinations to try to detect those cows suffering (or at risk of suffering) concurrent disease.

A well-constructed and implemented programme executed through well-trained staff can work very successfully. It is, however, not uncommon to find them diluted down to simply rectal temperatures in UK herds. Unfortunately, rectal temper-

Table 2. Risk factors for metritis²⁵

Risk factor	Odds ratio
Stillbirth	1.5
Assisted calving	2.1
Retained fetal membranes	6.0
Clinical ketosis	1.7

Table 3. Risk factors for left displaced abomasum²⁵

Risk factor	Odds ratio
Clinical hypocalcaemia	2.3
Clinical ketosis	13.8
Assisted calving	2.3

ature is often greater than the accepted normal range during the first 10 days after parturition^{23,24} and this occurs in response to the detection of proinflammatory cytokines (IL-1, IL-6, and TNF α) by receptors stimulating a coordinated neural response in the hypothalamus and brainstem to reset the thermostatic set point for body temperature²³.

A more risk-based approach can be taken to selecting cows for further examination by identifying those at higher risk of periparturient disease. These might include those that suffered an assisted calving, stillbirth, retained fetal membranes or clinical hypocalcaemia/ketosis, or gave birth to

Table 1. Suggested stocking densities for straw-bedded yards²²

Straw-bedded lying area	1.25m ² / 1,000L/cow
Loafing area*	3m ² /cow

*Loafing areas are defined as non-lying, non-passageway, non-feeding areas




Figure 2. Suggested cubicle dimensions for Holstein cows, adapted from Dairy Housing: a Best Practice Guide²¹.




Figure 3. Prevalence of uterine disorders in dairy cattle relative to calving²⁷

Summary

Maximising DMI through ration preparation and presentation, together with good environmental management, should promote a good lactation and reduce the incidence of related gateway diseases. A risk of successful lactation can be further increased by the prompt detection and appropriate treatment of periparturient diseases.

References

- Rastani R et al (2005). Reducing dry period length to simplify feeding transition cows: milk production, energy balance, and metabolic profiles, *Journal of Dairy Science* **88**(3): 1,004-1,014.
- NRC (2005). *Mineral Tolerance of Animals* (2nd edn), The National Academies Press.
- Suriyasathaporn W et al (2000). Hyperketonemia and the impairment of udder defense: a review, *Veterinary Research* **31**(4): 397-412.
- Rehage J et al (1996). Post surgical convalescence of dairy cows with left abomasal displacement in relation to fatty liver, *Schweizer Archiv Fur Tierheilkunde* **138**(7): 361-368.
- Wada Y, Muto M and Matsuura K (1995). Prognosis of cows with displaced abomasum and fatty infiltration of the liver, *Journal of the Japan Veterinary Medical Association* **48**(6): 387-390.
- Duffield T (1997). *Effects of a monesin controlled release capsule on energy metabolism, health and production in lactating dairy cattle*, University of Guelph: Ontario.
- Erb HN et al (1985). Path model of reproductive disorders and performance, milk fever, mastitis, milk yield, and culling in Holstein cows, *Journal of Dairy Science* **68**(12): 3,337-3,349.
- Butler WR and Smith RD (1989). Interrelationships between energy balance and postpartum reproductive function in dairy cattle, *Journal of Dairy Science* **72**(3): 767-783.
- Mayne C et al (2002). An investigation of fertility performance in dairy herds in Northern Ireland, *The Veterinary Record* **150**: 707-713.
- Hill A, Reid I, and Collins R (1985). Influence of liver fat on experimental *Escherichia coli* mastitis in periparturient cows, *The Veterinary Record* **117**(21): 549-551.
- Morrow DA et al (1979). Clinical investigation of a dairy-herd with the fat cow syndrome, *Journal of the American Veterinary Medical Association* **174**(2): 161-167.
- Houe H et al (2001). Milk fever and subclinical hypocalcaemia – an evaluation of parameters on incidence risk, diagnosis, risk factors and biological effects as input for a decision support system for

- Heinonen K et al (1987). The effect of mild fat infiltration in the liver on the fertility of Finnish Ayrshire cows, *Acta Veterinaria Scandinavica* **28**(2): 151-155.
- Duffield T (2000). Subclinical ketosis in lactating dairy cattle, *The Veterinary Clinics of North America: Food Animal Practice* **16**(2): 231-253.
- Heinonen K et al (1987). The effect of mild fat infiltration in the liver on the fertility of Finnish Ayrshire cows, *Acta Veterinaria Scandinavica* **28**(2): 151-155.
- Curtis C et al (1983). Association of parturient hypocalcemia with eight periparturient disorders in Holstein cows, *Journal of the American Veterinary Medical Association* **183**(5): 559-561.
- DeVries TJ, von Keyserlingk MA and Weary DM (2004). Effect of feeding space on the inter-cow distance, aggression, and feeding behavior of free-stall housed lactating dairy cows, *Journal of Dairy Science* **87**(5): 1,432-1,438.
- Cook N (2008). Pen moves and facility designs to maximize transition cow health and productivity, *Proceedings of the Penn*

- State Nutrition Conference.*
- Coonen J et al (2011). Short communication: effect of a stable pen management strategy for precalving cows on dry matter intake, plasma nonesterified fatty acid levels, and milk production, *Journal of Dairy Science* **94**(5): 2,413-2,417.
- Grant R (2007). Effect of stocking density on cow comfort, health and productivity, *Proceedings of the Cornell Nutrition Conference for Feed Manufacturers*, Syracuse, New York.
- AHDB (2012). *Dairy Housing: a Best Practice Guide*, Agriculture and Horticulture Development Board – Dairy.
- Green MJ et al (2007). National intervention study of mastitis control in dairy herds in England and Wales, *The Veterinary Record* **160**(9): 287-293.
- Saper C and Breder C (1994). The neurologic basis of fever, *New England Journal of Medicine* **330**(26): 1,880-1,886.
- Wagner S, Schimek D and Chen F (2008). Body temperature and white blood cell count in postpartum dairy cows, *The Bovine Practitioner* **42**(1): 18-26.

- Correa MT, Erb H and Scarlett J (1993). Path analysis for seven postpartum disorders of Holstein cows, *Journal of Dairy Science* **76**(5): 1,305-1,312.
- Sheldon IM (2004). The postpartum uterus, *The Veterinary Clinics of North America: Food Animal Practice* **20**(3): 569-591.
- Sheldon IM et al (2006). Defining postpartum uterine disease in cattle, *Theriogenology* **65**(8): 1,516-1,530.
- Sheldon IM, Rycroft A, and Zhou C (2004). Association of postpartum pyrexia and uterine bacterial infection in dairy cattle, *The Veterinary Record* **154**(10): 289-293.
- Sheldon IM et al (2009). Defining postpartum uterine disease and the mechanisms of infection and immunity in the female reproductive tract in cattle, *Biology of Reproduction* **81**(6): 1,025-1,032.
- Sheldon IM et al (2004). Minimum inhibitory concentrations of some antimicrobial drugs against bacteria causing uterine infections in cattle, *The Veterinary Record* **155**(13): 383-387.
- Noakes D (2001). Puerperal metritis. In Noakes D et al (eds), *Arthur's Veterinary Reproduction and Obstetrics*, Saunders.

JIM WILLSHIRE



is a partner of Endell Veterinary Group, based in Salisbury, and works in the farm animal department. In 2012 he was awarded the RCVS Diploma in Cattle Health and Production, for which he now examines. He sits on the DairyUK antibiotics working group, the research and development committee of Centre for Innovation Excellence in Livestock, representing XLVets, and is an external expert for Responsible Use of Medicines in Agriculture. Jim is also undertaking an MBA through the Open University.

32. British Veterinary Association (2009). *Responsible use of antimicrobials in veterinary practice: the 8-point plan* (<http://bit.ly/1rtYAAH>).

33. World Health Organization (2009). *Critically Important Antimicrobials for Human Medicine* (2nd revision; <http://bit.ly/1StPMk7>).

Table 4. Grading system for metritis²⁹

Grade 1 metritis	Cows with an abnormally enlarged uterus and a purulent uterine discharge without any systemic signs of ill health
Grade 2 metritis	Animals with additional signs of systemic illness such as decreased milk yield, dullness or pyrexia
Grade 3 metritis	Animals with signs of toxæmia, such as inappetence, cold extremities, depression or collapse